

2016

Sudden death syndrome – A growing threat of losses in soybeans

Shrishail S. Navi

Iowa State University, ssnavi@iastate.edu

X. B. Yang

Iowa State University, xbyang@iastate.edu

Follow this and additional works at: http://lib.dr.iastate.edu/plantpath_pubs



Part of the [Agricultural Science Commons](#), [Agriculture Commons](#), and the [Plant Pathology Commons](#)

The complete bibliographic information for this item can be found at http://lib.dr.iastate.edu/plantpath_pubs/104. For information on how to cite this item, please visit <http://lib.dr.iastate.edu/howtocite.html>.

This Article is brought to you for free and open access by the Plant Pathology and Microbiology at Iowa State University Digital Repository. It has been accepted for inclusion in Plant Pathology and Microbiology Publications by an authorized administrator of Iowa State University Digital Repository. For more information, please contact digirep@iastate.edu.

Sudden death syndrome – A growing threat of losses in soybeans

Shrishail S. Navi* and X.B. Yang

Address: 351 Bessey Hall, Department of Plant Pathology and Microbiology, Iowa State University, Ames, Iowa, USA

***Correspondence:** Shrishail S. Navi. Email: ssnavi@iastate.edu

Manuscript number: PAVSNNR-D-16-00098.

Received: 31 August 2016

Accepted: 12 October 2016

doi: 10.1079/PAVSNNR201611039

Article available online: <http://www.cabi.org/cabreviews/>

Abstract

Sudden death syndrome (SDS) caused by *Fusarium virguliforme* is one of the major yield-limiting soil borne diseases of soybean (*Glycine max*). The SDS has been reported from 21 U.S. states and is known to occur in Africa, North America, and South America. In the U.S. the losses due to SDS was estimated at \$3.06 billion for a period from 1988 to 2010. Since 1983, several management approaches have been investigated to reduce SDS and yet, continued efforts are necessary to develop long term disease management programs and to sustain disease below economic threshold levels. Integrating available control measures is an option, but adaptability and real-world assessments are equally important. Support of several funding agencies to better understand the disease in identifying suitable control measures to reduce yield losses in commercial cultivations has been indispensable in accomplishing these goals. In spite of sustained efforts, SDS continued to spread within the U.S. and reported in seven other countries since its first report in 1971. Comprehensive reviews have previously been published on this disease by Roy *et al.* [98], Leandro *et al.* [58], and Hartman *et al.* [32]. In this review, updated information on geographic distribution and economic significance of SDS, epidemiology, factors affecting SDS, and management options for SDS including screening techniques have been compiled. Also, discussed significant gaps in use of plant, fungi and bacteria based biocontrol agents in addressing management of SDS.

Keywords: Soybean sudden death syndrome, SDS, *Fusarium virguliforme*, biocontrol, global reports, yield losses due to SDS, epidemiological factors, greenhouse and field screening.

Review methodology: The following keywords were used in different combinations in searches on google, google scholar and CAB direct; first report(s) of soybean sudden death syndrome; yield losses due to soybean SDS; historical soybean prices per bushel or metric ton; factors affecting soybean SDS; management of soybean SDS; greenhouse and field screening techniques for SDS; biocontrol of soybean SDS; fungicide seed treatments; bio-fertilizer; host range; *Fusarium brasiliense*; *F. crassistipitatum*; *F. solani* f. sp. *glycines*; *F. tucumaniae*, and *F. virguliforme*; interaction between cyst nematode or *Heterodera glycines* on soybean SDS; back references cited in published articles, graduate theses, and unpublished data of several authors.

An overview of soybean

Taxonomy of the *Glycine* spp., and domestication [45], origin, history, and uses of soybean [27, 45, 61], history and growth of soybean plants including germplasm collections, utilization, plant improvement, growth and development [44] have been well documented. Soybean [*Glycine max* (L.) Merr.] is the leading oilseed crop produced and consumed in the world with the crop grown in 70 countries with an annual production of 268 million metric tons (mmt). Average world soybean production has increased from 115 mmt in 1993 to 308 mmt in 2014 and soybean prices (USD/ton) in top five producers (Argentina, Brazil, China mainland, India, and United States) fluctuated significantly between 1993 and 2014 [24]. According to Hymowitz *et al.* [44] top producers of soybean are the United States (31% of the world's total), Brazil (31%), Argentina (19%), China (5%), India (4%), Paraguay (3%) and Canada (2%). As of August 2016, USDA global production analysis projection of 2016/2017 world soybean production was 330.41 mmt, an increase of 17.74 mmt or a 5.67% compared with previous year [pecad.fas.usda.gov]. According to USDA Agricultural Projections to 2025, world trade is projected to increase in soybeans by 22%, soybean meal by 20%, and soybean oil by 30% [ers.usda.gov].

Global occurrence of sudden death syndrome

More than 200 plant pathogens affect soybean worldwide, of which at least 35 have been reported economically important [37]. Details of diseases in some of the major soybean producer countries such as Argentina, Brazil, Canada, China, India, Japan, and the United States have been compiled

by Hartman *et al.* [36]. Sudden death syndrome of soybean, caused by an Ascomycete fungus *Fusarium virguliforme* O'Donnell & T. Aoki in North America [7], and South Africa [125], *F. brasiliense* T. Aoki & O'Donnell, *F. crassistipitatum* Scandiani *et al.*, *F. cuneirostrum* O'Donnell & T. Aoki, *F. tucumaniae* T. Aoki *et al.*, and *F. virguliforme*, in South America [6–8, 90] is an economically significant soil-borne disease, and a risk to many soybean production areas worldwide. The disease was first observed by H.J. Walters in Arkansas, United States in 1971 [100], but it was only in 1983 that the disease was named as sudden death syndrome (SDS) of unknown etiology [40]. As of this review, the disease has been reported in three continents, North America (Canada and United States), South America (Argentina, Bolivia, Brazil, Paraguay, and Uruguay), and Africa (South Africa), and within the United States 21 states (Fig. 1). Also, *F. virguliforme* was isolated from soil in Malaysia [16], however, occurrence of SDS on soybean is yet to be recorded in that country. Similarly, *F. solani* f. sp. *glycines* has been reported in India not as SDS-causing pathogen but as a soybean wilt causing pathogen [Personal communication with Drs. Shamarao Jahagirdar, AICRP on Soybean, University of Agricultural Sciences, Dharwad, Karnataka, India, July 2016 and Rajesh K. Varma, ICAR-Indian Institute of Soybean Research, Indore, Madhya Pradesh, India, July 2016]. Also, in Japan, *F. azukicola* T. Aoki, Suga, F. Tanaka, Scandiani & O'Donnell, was able to induce root rot and typical SDS symptoms in soybean [9]. Whether the pathogens causing SDS have been introduced in to new regions or if they have been present in the soil in production fields of these countries for some time without being detected is unknown. After the SDS was detected in Iowa [150], Scherm and Yang [114] developed a risk assessment model to determine potential geographic range of development of this disease, which was considered a southern disease by then, in North America. They predicted potential northward development of SDS and cooler production regions in the North Central United States to have higher risk than southern soybean production regions. Eighteen years after their publication, occurrence and distribution of SDS confirmed accuracy of their predictions [114]. Future reports of SDS in India, Japan and Malaysia on soybeans would be an interesting development of this disease.

Economic significance of SDS

Historic yield losses due to SDS in the United States compiled from various sources [21, 35, 53, 96, 116, 132–134, 136–141] ranged from 0.06 mmt in 1988 to 1.91 mmt in 2010 with the highest yield loss of 2.10 mmt recorded in 2000. The corresponding economic losses due to SDS varied from \$15.7 million in 1988 to \$669.2 million in 2010 (Fig. 2). The economic losses were based on NASS-USDA soybean prices at \$7.5/bushel (\$275.6/metric ton) in 1988 and \$9.6/bushel (\$352.7/metric ton) in 2010 (Fig. 2). Though the estimated yield loss of 2.10 mmt in 2000 was highest compared to yield loss of 1.91 mmt in 2010, due to differences in soybean prices the economic loss in 2000 was \$342.7 million (\$4.5/bushel or \$163.5/metric ton), which is almost half the economic loss recorded in 2010 at \$669.2 million. On the contrary, the soybean prices in 2008 were highest (\$385.1/metric ton) compared with other years between 1988 and 2010, the economic loss was estimated at \$209.6 million due to less yield losses of only 0.54 mmt in 2008. An estimated average yield loss due to SDS from 1988 to 2010 was 0.56 mmt (range 0.03–2.10 mmt) and corresponding average economic loss was \$133.3 million (range \$8.1–669.2 million) at an average soybean price at \$233.24/metric ton (range \$152.5–385.1/metric ton). Yield losses due to SDS vary widely depending on planting dates [39, 123, 135], maturity groups, cultivar selection [123], climatic and environmental trends during the growing season [57], glyphosate spray and many more dynamics as outlined in factors affecting SDS. Also, economic losses vary depending on the market price of soybean in a given year and primary factors affecting the market price according to USDA Economic Research Service include, population and income growth, demand for livestock products, as well as export import policies. In this review, the yield and economic losses due to SDS in Canada, South American countries and South Africa were not calculated because of non-availability of historical data.

Symptoms of SDS

Soybean plants infected with SDS show symptoms both during seedling (Fig. 3A-E) and the most dramatic symptoms in reproductive growth stages (Fig. 3F-K) similar to Roy *et al.* [98]. Characteristic symptoms include, appearance of small yellow flecks or chlorotic spots that coalesce to cause interveinal chlorosis. As the disease progresses, interveinal chlorosis turns into interveinal necrosis and the affected leaves twist and curl giving puckering and mottling appearance that defoliate prematurely with leaf petiole remaining intact on the stem. Flowers and

Pods abort and developing pods may not entirely fill. Symptomatic plants exhibit blue-stained fungal sporodochia on taproot (Fig. 3K). Factors that favor symptom development are cool, moist conditions early in the growing season leading to higher disease incidence. Also, early planting, high rainfall and/or low-lying, poorly drained or compacted areas of the fields are other important factors that lead to higher incidence. Above-ground symptoms are caused mainly by a toxin produced by the fungus and translocated through the plant. The disease severity varies from field to field depending on the above conditions and field history.

Infection process and survival of *F. virguliforme*

The occurrence of *F. virguliforme* infection early in the season is the result of colonization in xylem tissue, which provides a pathway for upward movement of phytotoxin that are essential for foliar symptom expression [83, 86] and lignin degradation by *F. virguliforme* may play an important role in the infection, colonization, and survival of the fungus in the root tissue [62, 63]. Navi and Yang [83] reported that, plants with SDS foliar symptoms showed both external and internal discolored taproots and basal stems, while plants with no foliar symptoms had only superficial external discoloration. Microtome sectioning of taproots of symptomatic plants revealed the presence of fungal structures in both xylem and phloem tissues, while plants with no foliar symptoms revealed fungal structures only in phloem tissue. Based on their findings Navi and Yang [83] suggested an effective and ineffective colonization zones of the fungus for symptomatic and asymptomatic plants, respectively (Fig. 4).

The SDS fungus survives between crops in soil and crop residue (corn and soybean) either as conidia or chlamydospore [84, 98], on decomposed soybean roots post-harvest [100, 101], inside soybean cyst nematode [67, 68] and in no-till fields [71, 100]. This pathogen can also reproduce on corn, wheat, ryegrass, pigweed, sugar beet, lambsquarters, canola, alfalfa, pinto bean, navy bean, white clover, red clover, pea, and Canadian milk vetch without causing symptoms on these hosts [55]. Researchers have been puzzled by the fact that SDS has now become more prevalent in the Corn Belt and there were many observations that severe SDS occurred after continuous corn. Recently, Navi and Yang [84] investigated effects of rotation with corn on *F. virguliforme* survival. Their results showed that corn residue, particularly coarse-ground corn kernels, harbor *F. virguliforme* significantly in the absence of soybeans. Also, studies conducted

by Abdelsamad *et al.* [1] with a 2-year rotation of corn and soybean showed greater SDS incidence and severity, and lower yield, compared to the 3-year corn-soybean-oat/red clover, and 4-year corn-soybean- oat/alfalfa-alfalfa rotations. Thus *F. virguliforme* remains a growing threat to global soybean production.

Factors affecting occurrence of SDS

There are numerous factors that affect SDS such as; effects of herbicides on metabolic activities and growth of host and pathogens and increase in soil-borne diseases as a result of herbicide-pathogen interaction [3], interaction between *F. virguliforme* and cyst nematodes [26, 99, 103, 143, 144], biotic and abiotic factors [98], irrigation and cyst nematodes [70], irrigation and soil compaction [20], tillage [117, 127, 128, 135], inoculum density [28], isolates and inoculum rate [31], genetic structure and variation in aggressiveness of *F. virguliforme* [66], soil variables including fertility parameters [105, 112], soil temperature [29, 113], water matric potential in soil [113], relation of sand content, soil pH, and soil nutrients [109], cultivars [39, 89, 101, 105, 127, 135], genetic architecture of soybean [156], planting date [39, 52, 135], crop rotation and crop sequences [93, 104, 142], root system [91], plant age [29, 30], early infection and xylem colonization [83, 86, 154], biochemical response of soybean roots [62], toxin production by pathogen [12], herbicides spray [22, 50, 77–79, 107, 108, 147], light [49], elevated atmospheric carbon dioxide and ozone [23], climatic and environmental trends [57], symptomatic and asymptomatic host-range [55], fungicide and cultivar interactions [51], and seed treatment and plant population [2, 52, 155].

Management options of SDS

The sudden occurrence and unpredictable nature of soybean SDS make the disease so far the most puzzling one to manage in soybean production. Development of the disease depends on several factors listed above. However, a number of approaches listed below either alone or by integration of several methods can help reduce SDS impact in commercial production. Management options of SDS to break the life cycle of the pathogen (Fig. 5) include delayed planting [39, 123, 135], planting SDS tolerant varieties [102, 108], fall tillage [135, 137], crop rotation [1, 104, 137, 142],

modifying agronomic practices like row spacing and seeding rate [123], fungicide seed treatments [52, 72, 82, 130, 131, 155], seed treatments with a combination of fungicides, systemic insecticides and biological(s) [2, 155], seed treatment with bacteria and fungi based biocontrol agents [43, 75, 87, 129], preplant or foliar applied potassium chloride with fungicides [88], cultural and biological control [4], exploring potential untapped resistance sources in perennial *Glycine* spp. to improve resistance in soybean [33] and recovering SDS infected plants [79] similar to sorghum and pearl millet [120, 121], plant resistance, variety selection, adjusting planting dates, crop rotation, seed treatment with bio-fungicide [149], identification of quantitative trait loci [145], genomic approaches to molecular breeding of resistance [46], integrated approaches [34], clean harvest of corn and soybean [84, 151] and genetic engineering along with other traditional management options may be needed as integrated approaches to manage SDS [32].

Recently, Navi and Yang [75] have summarized HeadsUp (a plant based biocontrol product) seed treatment or foliar spray studies of 11 years starting in 2005 covering six states in North Central regions of the United States. Results showed that HeadsUp has effectively suppressed SDS and white mold (*Sclerotinia sclerotiorum* (Lib.) de Bary) and improved an average yield of 0.142 mt/ha (range -0.830 to 1.095 mt/ha) compared with controls (Fig. 6). Compared with control, the proportions of yield response to treatments were 62% increase, 9% no change, and 29% decrease in 68 field trials. There is a slightly larger number of low responsive trials for HeadsUp compared with synthetic chemicals, which is often seen in field trials with biological control products. However, the low cost and environmental friendly make HeadsUp a good candidate for soybean disease management.

Resistance screening in greenhouse conditions

Several greenhouse methods for host resistant screening that are previously reported in the literature that have been successful in identifying soybean resistant to SDS. These include dip inoculation of sprouted seed in *F. virguliforme* spore suspension and planting in greenhouse mix [80, 81, 85], temperature-controlled water bath method [38], exposing cut stem to cell free culture filtrate [42], molecular marker-assisted breeding for resistance [60], toothpick inoculation and soil infestation methods [111], planting seeds in a mixture of 1 part inoculum with 20 part soil [64], using cones filled with steam-treated soil mix (2:1 sand: soil) and topped with 3 g of fungus

infested white sorghum grains [25]. Navi and Yang (unpublished), modified greenhouse screening technique [80, 81, 85] to identify resistance sources in germplasm and explore efficacy tests of seed treatments with biocontrol agents (BCA) and or chemicals against SDS (Fig. 7). The modified method of Navi and Yang [unpublished] in brief is as follows; White sorghum grains infested with *F. virguliforme* [84] were sprinkled into plastic tub containing potting mixture (two parts of steam-sterilized sand and one part black soil) at 1% of the potting mixture (Fig. 7A), and the inoculum was mixed with the potting mixture (Fig. 7B). White foam cups (237 ml, Dart Container of Michigan LLC, 500 Hogsback Rd., Mason, MI 48854) were filled-up with the mixture, and the planting spots were marked at equidistant (1.5-cm apart) and at uniform depth of 2-cm (Fig. 7C) with hand held wooden planter (Fig. 7D) fabricated at General Services workshop facility, Iowa State University. Placed 5-7 seeds (commercially untreated seeds that were treated with either BCA spore suspension or fungicide at 5 ml/kg of seed) in marked spots (Fig. 7E) and pressed the seed to uniform depth of 2-cm with the planter. The seed was covered with additional potting mixture and was compacted it (Fig. 7F) with hand held wooden compactor (Fig. 7G) and watered the cups to the saturation point. Cups were incubated on greenhouse benches in 16-h light under 400W E-Ballast, Metal Halide type M59 bulb, (120v/HPS/MH, Digital Sunburst, Hydrofarm Inc. 2249 S. McDowell Ext., Petaluma, CA, 94954). During the incubation, plants were watered twice daily. Two weeks after planting, stand count and initial SDS counts were recorded. Subsequently, 30 and 45 days after planting, SDS infected plants count and SDS plants showing production of symptomless leaves that previously had one or more unifoliate and or trifoliate leaves with SDS symptoms and their severities were recorded. Authors are using this method from 2010 till date to evaluate seed treatment industry protocols and in academic research.

Resistance screening in field conditions

Identifying resistance to SDS remains a major effort for soybean industry because variety selection is the most effective source for producers in SDS risk management. Resistance breeding has made progress over the past 20 years with a limited success due to the fact that SDS resistance is controlled by multiple genes, which makes resistance screening for a large number of entries difficult. Currently, selections and evaluations for resistance to SDS largely depends on both greenhouse and field evaluations. Every season, breeding lines are evaluated in fields either with

history of SDS or in fields inoculated with *F. virguliforme* fermented on oats a method initially developed by Yang *et al.* [148]. In this method, SDS incidence and severity increased with an increased ratio of oat to soybean seeds, reaching a maximum level at about 4:1. Subsequently, this method was modified in 2006 [20] for establishing field nurseries and in 2008 [Navi and Yang, unpublished] to assess efficacy of biological and chemical seed treatments or seed treatment plus foliar sprays. The modified method of Navi and Yang [unpublished] in brief is as follows; increased *F. virguliforme* on white sorghum grain [84], sampled separately a known number of soybean seed at 10 seed/linear foot (76.2-cm) in 8.5 × 20.3-cm envelopes and the *F. virguliforme* fermented sorghum grain at 5cc/linear foot. The seeds of individual treatment and the inoculum were then placed into a cone plot planter (2- or 4-rows). Planted soybean seed along with the fermented sorghum grain using ALMACO 4-Row SeedPro Precision Vacuum planter with an automatic cable winding trip system. Applied post-emergence herbicide (if required) prior to setting up rainbird overhead sprinklers. Run the sprinklers on non-rainy days from vegetative growth stage-4 (four unfolded trifoliolate leaves) to reproductive growth stage-6 (full seed setting) to provide either irrigation water or rainfall a total of 2.5 cm/week. Recorded stand count in vegetative growth stages at 15 and 30 days after planting and SDS plants count and severities both in vegetative (Fig. 3A-D) and reproductive growth stages (Fig. 3F-J) at regular intervals up to R8. This method has an increased consistency of SDS symptom expression across replications in testing efficacy of biological and chemical seed treatment products and or the foliar sprays against SDS and is extensively used by the authors to test industry protocols. Also, the method is in use to test advanced breeding lines with different maturity groups [19].

Conclusion and future research opportunities

Understanding the nature of sudden death of a soybean plant colonized by SDS fungus and occurrence of sudden blight of a soybean field due to SDS so far remained the major challenge in developing effective management measures. Development of a SDS resistance screening method which can handle a large number of entries in a short period of time remains one of the major challenges for SDS research. Currently, the seed treatment with ILeVo appears promising, although foliar symptoms do occur if the variety is susceptible to SDS. Information technology enabled us to study this disease using a non-traditional approach, which may provide new insight

on the sudden occurrence of this disease. Yang *et al.* [146] showed SDS to be a model system in the study of satellite remote sensing for disease detection. This is the first reported case of occurrence of a plant disease that can be seen and identified from satellite due to its unique nature.

Based on currently available literature search, several possible scenarios have been discussed to improve soybean SDS management. However, we have observed that there is a significant gap in the use of fungi or bacteria based biocontrol agents (BCA) in management of SDS. Therefore, much needed emphasis is essential on use of BCA that have strong mycoparasitism and antagonism characters. Also, BCA seed treatment with bioAPT [87] a microbial carrier powder (American Peat Technology, LLC, Aitkin, MN, U.S.A.) or without the carrier [43, 129], application of BCA along with pre-planting fertilizers and herbicides, and or foliar spray need to be standardized to have suitable synergism and compatibility. Although, *Fusarium* spp. causing SDS are not seed borne except one report [10], but seed treatments with BCA provide mycoparasitism potentials of the target pathogen in the vicinity of seed germination zone by which early infection and subsequent toxin production could be minimized. In addition, application of BCA in any of the arrangements stated above should have long term benefits as most fungi or bacteria based BCA increase their population much faster than the target pathogen on crop residue and if conditions are unfavorable they survive in soil and residue by producing chlamydospores. If the BCA has potential to mycoparasitize hyphae, conidia or chlamydospores of the target pathogen, it is predictable to have an added advantage.

The activity of BCA is mainly attributed to various anti-microbial/antagonistic compounds they produce [13, 15, 41], competition for nutrition and space [14, 59, 122], antibiosis [115], antagonism effects [71], and mycoparasitism capabilities [17, 48, 97]. Full exploitation of the BCA potential could provide growth enhancement of domestic plants, green house plants, and agricultural crops [65, 126]. Becker Underwood Inc. (now with BASF chemical company) has developed bacterium based HiStick N/T (currently Nodulator N/T) a peat-based inoculant and disease protection seed treatment product in soybean to suppress *Rhizoctonia* spp. and *Fusarium* spp. (BASF.com). Also, Organic Materials Review Institute (omri.org) has developed *Trichoderma* spp. based soil amendment and bio-fertilizer (Custom GP) to improve overall conditions of the soil aimed at healthier roots and stronger plant growth. Recently, a team led by scientist Aradhana Mishra, National Botanical Research Institute, one of the constituent research institutes of the Council of Scientific and Industrial Research, New Delhi, India developed

Trichoderma spp. based bio-fertilizer to mitigate greenhouse gases and to repair damaged root tissues of various agriculture crops, thereby increasing productivity (article appeared in Times of India, Indian National Newspaper, June 6, 2016). Wide-ranging commercial BCA registered under the United States Environmental Protection Agency, *Trichoderma* spp. based biofungicide agents and agriculture products [106, 126] and other commercially available biocontrol products used against several plant diseases [76] can be explored in management of soybean SDS.

Acknowledgements

We thank Dr. Krishna V. Subbarao, Professor, Department of Plant Pathology, University of California-Davis, U.S.A. and Dr. A.H. Rajasab, Professor and Vice-chancellor, Tumkur University, Karnataka, India for critical review of the manuscript.

Disclaimer

Some of the products indicated in this review do not imply endorsement of one company over another, nor was discrimination intended against any similar products.

References

1. Abdelsamad N, Mbofung GC, Robertson AE, Liebman M, Leandro LF. Long-term crop rotations suppress soybean sudden death syndrome in Iowa. *Phytopathology*. 2012;102(7):S4.1.
2. Adee EA. Effects of seed treatment on sudden death syndrome symptoms and soybean yield. *Kansas Agricultural Experiment Station Research Reports*. 2015;1:3.
3. Altman J, Rovira AD. Herbicide-pathogen interactions in soil-borne root diseases. *Canadian Journal of Plant Pathology*. 1989;11:166–72.
4. Anderson TR, Gnanamanickam SS. Biological control of soybean diseases. *Biological Control of Crop Diseases*. 2002;3:191.
5. Anderson TR, Tenuta AU. First report of *Fusarium solani* f. sp. *glycines* causing sudden death syndrome of soybean in Canada. *Plant Disease*. 1998;82:448.

6. Aoki T, O'Donnell K, Scandiani MM. Sudden death syndrome of soybean in South America is caused by four species of *Fusarium*: *Fusarium brasiliense* sp. nov., *F. cuneirostrum* sp. nov., *F. tucumaniae*, and *F. virguliforme*. *Mycoscience*. 2005;46:162–83.
7. Aoki T, O'Donnell K, Homma Y, Lattanzi AR. Sudden-death syndrome of soybean is caused by two morphologically and phylogenetically distinct species within the *Fusarium solani* species complex—*F. virguliforme* in North America and *F. tucumaniae* in South America. *Mycologia*. 2003;95:660–84.
8. Aoki T, Scandiani MM, O'Donnell K. Phenotypic, molecular phylogenetic, and pathogenetic characterization of *Fusarium crassistipitatum* sp. nov., a novel soybean sudden death syndrome pathogen from Argentina and Brazil. *Mycoscience*. 2012;53:167–86.
9. Aoki T, Tanaka F, Suga H, Hyakumachi M, Scandiani MM, O'Donnell K. *Fusarium azukicola* sp. nov., an exotic azuki bean root-rot pathogen in Hokkaido, Japan. *Mycologia*. 2012 Sep 1;104(5):1068–84.
10. Balardin CR, Celmer AF, Costa EC, Meneghetti RC, Balardin RS. Possible transmission of *Fusarium solani* f. sp. *glycines*, causal agent of sudden death syndrome, through soybean seed. *Fitopatologia Brasileira*. 2005;30:574–81.
11. Bernstein ER, Atallah ZK, Koval NC, Hudelson BD, Grau CR. First report of sudden death syndrome of soybean in Wisconsin. *Plant Disease*. 2007;91:1201.
12. Brar HK, Swaminathan S, Bhattacharyya MK. The *Fusarium virguliforme* toxin FvTox1 causes foliar sudden death syndrome-like symptoms in soybean. *Molecular Plant-microbe Interactions*. 2011;24:1179–88.
13. Brozová J. Mycoparasitic fungi *Trichoderma* spp. in plant protection. A review. *Plant Protection Science-UZPI (Czech Republic)*. 2004.
14. Celar F. Competition for ammonium and nitrate forms of nitrogen between some phytopathogenic and antagonistic soil fungi. *Biological Control*. 2003;28:19–24.
15. Chaube HS, Mishra DS, Varshney S, Singh SU. Biocontrol of plant pathogens by fungal antagonists: Historical background, present status and future prospects. *Annual Review of Plant Pathology*. 2003;2:1–42.
16. Chehri K, Salleh B, Zakaria L. *Fusarium virguliforme*, a soybean sudden death syndrome fungus in Malaysian soil. *Australasian Plant Disease Notes*. 2014;9:1–7.

17. Chet I, Harman GE, Baker R. *Trichoderma hamatum*: Its hyphal interactions with *Rhizoctonia solani* and *Pythium* spp. Microbial Ecology. 1981;7:29–38.
18. Chilvers MI, Brown-Rytlewski DE. First report and confirmed distribution of soybean sudden death syndrome caused by *Fusarium virguliforme* in Southern Michigan. Plant Disease. 2010;94:1164.
19. Cianzio SR, Gebhart GD, Lundeen P. Soybean Sudden Death Syndrome Field Screening. 2008. Iowa State University, Northern Research and Demonstration Farm, ISRF08-22. Ames, Iowa, USA.
20. de Farias Neto AL, Hartman GL, Pedersen WL, Li S, Bollero GA, Diers BW. Irrigation and inoculation treatments that increase the severity of soybean sudden death syndrome in the field. Crop Science. 2006;46:2547–54.
21. Doupnik BJr. Soybean production and disease loss estimates for north central United States from 1989 to 1991. Plant Disease. 1993;77:1170-71.
22. Duke SO, Lydon J, Koskinen WC, Moorman TB, Chaney RL, Hammerschmidt R. Glyphosate effects on plant mineral nutrition, crop rhizosphere microbiota, and plant disease in glyphosate-resistant crops. Journal of Agricultural and Food Chemistry. 2012;60:10375–97.
23. Eastburn DM, Degennaro MM, Delucia EH, Dermody O, Mcelrone AJ. Elevated atmospheric carbon dioxide and ozone alter soybean diseases at SoyFACE. Global Change Biology. 2010;16:320–30.
24. FAOSTAT <http://faostat3.fao.org> Food and Agriculture Organization of the United Nations Statistics Division.
25. Farias Neto AL, Schmidt M, Hartman GL, Shuxian L, Diers BW. Inoculation methods under greenhouse conditions for evaluating soybean resistance to sudden death syndrome. Pesquisa Agropecuária Brasileira. 2008;43:1475–82.
26. Gao X, Jackson TA, Hartman GL, Niblack TL. Interactions between the soybean cyst nematode and *Fusarium solani* f. sp. *glycines* based on greenhouse factorial experiments. Phytopathology. 2006;96:1409–15.
27. Gibson L, Benson G. Origin, History, and Uses of Soybean (*Glycine max*). Iowa State University, Department of Agronomy, March. 2005, Ames, Iowa, USA.

28. Gongora-Canul C, Nutter Jr FW, Leandro LF. Temporal dynamics of root and foliar severity of soybean sudden death syndrome at different inoculum densities. *European Journal of Plant Pathology*. 2012;132:71–9.
29. Gongora-Canul CC, Leandro LF. Effect of soil temperature and plant age at time of inoculation on progress of root rot and foliar symptoms of soybean sudden death syndrome. *Plant Disease*. 2011;95:436–40.
30. Gongora-Canul CC, Leandro LF. Plant age affects root infection and development of foliar symptoms of soybean sudden death syndrome. *Plant Disease*. 2011a;95:242–7.
31. Gray LE, Achenbach LA. Severity of foliar symptoms and root and crown rot of soybean inoculated with various isolates and inoculum rates of *Fusarium solani*. *Plant Disease*. 1996;80:1197–9.
32. Hartman GL, Chang HX, Leandro LF. Research advances and management of soybean sudden death syndrome. *Crop Protection*. 2015;73:60–6.
33. Hartman GL, Gardner ME, Hymowitz T, Naidoo GC. Evaluation of perennial species for resistance to soybean fungal pathogens that cause Sclerotinia stem rot and sudden death syndrome. *Crop Science*. 2000;40:545–9.
34. Hartman GL, Hill CB. Diseases of soybean and their management. 2010. 276. In Singh G, editor. *The soybean: botany, production and uses*. CABI. 2010.
35. Hartman GL, Noel GR, Gray LE. Occurrence of soybean sudden death syndrome in east-central Illinois and associated yield losses. *Plant Disease*. 1995;79:314–8.
36. Hartman GL, Rupe JC, Sikora EJ, Domier LL, Davis JA, Steffey KL. Soybean diseases in Argentina, Brazil, Canada, China, India and Japan. 2015a; Pages 5-16. In: *Compendium of soybean diseases and pests*, fifth edition. The American Phytopathology, 3340 Pilot knob road, St. Paul, MN, USA, 201pp. Edited by Glen L. Hartman, John C. Rupe, Edward J. Sikora, Leslie L. Domier, Kevin L. Steffey, and Jeffrey A. Davis
37. Hartman GL. Worldwide importance of soybean pathogens and pests. 2015. Pages 4-5. In: *Compendium of soybean diseases and pests*, fifth edition. The American Phytopathology, 3340 Pilot knob road, St. Paul, MN, USA, 201pp. Edited by Glen L. Hartman, John C. Rupe, Edward J. Sikora, Leslie L. Domier, Kevin L. Steffey, and Jeffrey A. Davis

38. Hashmi RY, Bond JP, Schmidt ME, Klein JH. A temperature-controlled water bath method for evaluating soybean reaction to sudden death syndrome (SDS). Online. Plant Health Progress. 2005. doi:10.1094/PHP-2005-0906-01-RS.
39. Hershman DE, Hendrix JW, Stuckey RE, Bachi PR, Henson G. Influence of planting date and cultivar on soybean sudden death syndrome in Kentucky. Plant Disease. 1990;74:761–6.
40. Hirrel MC. Sudden death syndrome of soybean: a disease of unknown etiology. Phytopathology 1983;73:501–2.
41. Howell CR. Cotton seedling preemergence damping-off incited by *Rhizopus oryzae* and *Pythium* spp. and its biological control with *Trichoderma* spp. Phytopathology. 2002;92:177–80.
42. Huang YH, Hartman GL. Reaction of selected soybean genotypes to isolates of *Fusarium solani* f. sp. *glycines* and their culture filtrates. Plant Disease. 1998;82:999–1002.
43. Huynh T, Yang X, Navi S, Li X. Reduction of soybean sudden death syndrome (*Fusarium virguliforme*) by seed treatment with *Bradyrhizobium japonicum* (Abstr). 2016. APS Annual Meeting, July 30 - Aug 3, Tampa, Florida, U.S.A.
<http://www.apsnet.org/meetings/annual/abstracts/pages/abstractdetail.aspx?MID=257>
44. Hymowitz T, Nelson R, Sinclair JB, Hartman GL. History and growth of the soybean plants. 2015. Pages 1-4. In: *Compendium of soybean diseases and pests*, fifth edition. The American Phytopathology, 3340 Pilot knob road, St. Paul, MN, U.S.A. Edited by Glen L. Hartman, John C. Rupe, Edward J. Sikora, Leslie L. Domier, Kevin L. Steffey, and Jeffrey A. Davis
45. Hymowitz T, Newell CA. Taxonomy of the genus *Glycine*, domestication and uses of soybeans. Economic Botany. 1981;35:272–88.
46. Iqbal J, Cryder K, Ahsan R, Triwitayakorn KT, Jamail A, Schulz J, et al. Genomic approaches to molecular breeding of resistance to soybean sudden death syndrome and cyst nematode in elite cultivars. In Proceedings VII World Soybean Research Conference, IV International Soybean Processing and Utilization Conference, III Congresso Brasileiro de Soja (Brazilian Soybean Congress), Foz do Iguassu, PR, Brazil, 29 February-5 March, 2004. 2004 (pp. 471-480). Brazilian Agricultural Research Corporation, National Soybean Research Center.

47. Jardine DJ, Rupe JC. First report of sudden death syndrome of soybeans caused by *Fusarium solani* in Kansas. Plant Disease. 1993;77:1264.
48. Jeffries P. Biology and ecology of mycoparasitism. Canadian Journal of Botany. 1995;73:1284–90.
49. Ji J, Scott MP, Bhattacharyya MK. Light is essential for degradation of ribulose-1, 5-bisphosphate carboxylase-oxygenase large subunit during sudden death syndrome development in soybean. Plant Biology. 2006;8:597–605.
50. Kandel YR, Bradley CA, Wise KA, Chilvers MI, Tenuta AU, Davis VM, et al. Effect of glyphosate application on sudden death syndrome of glyphosate-resistant soybean under field conditions. Plant Disease. 2015;99:347–54.
51. Kandel YR, Wise KA, Bradley CA, Chilvers MI, Tenuta AU, Mueller DS. Fungicide and cultivar effects on sudden death syndrome and yield of soybean. Plant Disease. 2016;100:1339–50.
52. Kandel YR, Wise KA, Bradley CA, Tenuta AU, Mueller DS. Effect of planting date, seed treatment, and cultivar on plant population, sudden death syndrome, and yield of soybean. Plant Disease. 2016a; 100:1735–43.
53. Koenning SR, Wrather JA. Suppression of soybean yield potential in the continental United States by plant diseases from 2006 to 2009. Plant Health Progress. 2010:1094.
54. Koenning SR. Soybean sudden death syndrome. Soybean Disease Information Note 7. 2002; <http://www.ces.ncsu.edu/depts/pp/notes/Soybean/soy007/soy007.htm>
55. Kolander TM, Bienapfl JC, Kurle JE, Malvick DK. Symptomatic and asymptomatic host range of *Fusarium virguliforme*, the causal agent of soybean sudden death syndrome. Plant Disease. 2012;96:1148–53.
56. Kurle JE, Gould SL, Lewandowski SM, Li S, Yang XB. First report of sudden death syndrome (*Fusarium solani* f. sp. *glycines*) of soybean in Minnesota. Plant Disease. 2003;87:449.
57. Leandro LF, Robertson A, Mueller DS, Yang XB. Climatic and environmental trends observed during epidemic and non-epidemic years of soybean sudden death syndrome in Iowa. Plant Health Progress. 2013. Online. Plant Health Progress doi:10.1094/PHP-2013-0529-01-RS.

58. Leandro LF, Tatalovic N, Luckew A. Soybean sudden death syndrome—advances in knowledge and disease management. *Plant Sciences Reviews*. 2012. 2012:215.
59. Lewis JA, Papavizas GC. Biocontrol of plant diseases: the approach for tomorrow. *Crop Protection*. 1991;10:95–105.
60. Lightfoot DA. Two Decades of molecular marker-assisted breeding for resistance to soybean sudden death syndrome. *Crop Science*. 2015;55:1460–84.
61. LiJuan Q, RuZhen C, Singh G. The origin and history of soybean. *The soybean: botany, production and uses*. 2010:1–23.
62. Lozovaya VV, Lygin AV, Li S, Hartman GL, Widholm JM. Biochemical response of soybean roots to *Fusarium solani* f. sp. *glycines* infection. *Crop Science*. 2004;44:819–26.
63. Lozovaya VV, Lygin AV, Zernova OV, Li S, Widholm JM, Hartman GL. Lignin degradation by *Fusarium solani* f. sp. *glycines*. *Plant Disease*. 2006;90:77–82.
64. Luckew AS, Cianzio SR, Leandro LF. Screening method for distinguishing soybean resistance to in resistant \times resistant crosses. *Crop Science*. 2012;52:2215–23.
65. Mastouri F, Björkman T, Harman GE. Seed treatment with *Trichoderma harzianum* alleviates biotic, abiotic, and physiological stresses in germinating seeds and seedlings. *Phytopathology*. 2010;100:1213–21.
66. Mbofung GY, Harrington TC, Steimel JT, Navi SS, Yang XB, Leandro LF. Genetic structure and variation in aggressiveness in *Fusarium virguliforme* in the Midwest United States. *Canadian Journal of Plant Pathology*. 2012;34:83–97.
67. McLean KS, Lawrence GW. Development of *Heterodera glycines* as affected by *Fusarium solani*, the causal agent of sudden death syndrome of soybean. *Journal of Nematology* 1995;27:70–7.
68. McLean KS, Lawrence GW. Interrelationship of *Heterodera glycines* and *Fusarium solani* in sudden death syndrome of soybean. *Journal of Nematology* 1993;25:434–9.
69. Meiring B, Dorrance A, Mills D. Sudden death syndrome of soybean, Fact Sheet Agricultural and Natural Resources, 2010;AC-44-10. <http://ohioline.osu.edu/ac-fact/pdf/0044.pdf>
70. Melgar J, Roy KW, Abney TS. Sudden death syndrome of soybean – etiology, symptomatology, and effects of irrigation and *Heterodera glycines* on incidence and severity under field conditions. *Canadian Journal of Botany*. 1994;72:1647–53.

71. Milanesi PM, Blume E, Muniz MF, Reiniger LR, Antonioli ZI, Junges E, et al. Detection of *Fusarium* spp. and *Trichoderma* spp. and antagonism of *Trichoderma* spp. in soybean under no-tillage. *Semina: Ciências Agrárias* (Londrina). 2013;34:3219–33.
72. Mueller TA, Knake RP, Riggs JL. Control of *Fusarium virguliforme* (sudden death syndrome) with a seed treatment. *Phytopathology*. 2011;101:S124.
73. Mulrooney RP, Gregory NF, Walker SD, Webster AM. First report of sudden death syndrome of soybean in Delaware and eastern shore of Maryland. *Plant Disease*. 2002;86:696.
74. Nakajima T, Mitsueda T, Charchar MD. First occurrence of sudden death syndrome of soybean in Brazil. *JARQ, Japan Agricultural Research Quarterly*. 1996;30:31–4.
75. Navi S, Yang X. Summary of 11-year field trials of using HeadsUp - a Saponin plant protectant to manage soybean diseases. 2016a. APS annual meeting, July 30 - Aug 3, Tampa, Florida, U.S.A.
<http://www.apsnet.org/meetings/annual/abstracts/pages/abstractdetail.aspx?MID=256>
76. Navi SS, Bandyopadhyay R. Biological control of fungal plant pathogens. 2002. Pages 354-365. In: *Plant Pathologist's Pocketbook*. JM Waller, JM Lenné and SJ Waller, eds. CABI Publishing, Wallingford, U.K.
77. Navi SS, Jing L, Yang XB. Effects of glyphosate application rates on soybean sudden death syndrome. *Phytopathology*. 2012;102:S4.
78. Navi SS, Jing L, Yang XB. Effects of glyphosate application rates and frequency on soybean sudden death syndrome. 2013. *Plant Pathology Presentations and Posters*. Paper 4.
http://lib.dr.iastate.edu/plantpath_conf/4
79. Navi SS, Jing L, Yang XB. Report of recovery of soybean from sudden death syndrome caused by *Fusarium virguliforme*. In *Phytopathology* 2014 Nov 1 (Vol. 104, No. 11, pp. 173–173). 3340 Pilot Knob Road, St Paul, MN 55121 USA: American Phytopathological Society.
80. Navi SS, Yang XB. A novel screening technique to identify resistance to soybean sudden death syndrome. *Phytopathology*. 2004;94(6). *Phytopathology* 94:S146. Publication no. P-2004-0024-SOA.
81. Navi SS, Yang XB. Dip inoculation technique to identify resistance to soybean sudden death syndrome. *Phytopathology*. 2003. 93:S65.

82. Navi SS, Yang XB. Effects of seed treatment on root diseases and yields of soybean. *Phytopathology*. 2010;100:S89.
83. Navi SS, Yang XB. Foliar symptom expression in association with early infection and xylem colonization by *Fusarium virguliforme* (formerly *F. solani* f. sp. *glycines*), the causal agent of soybean sudden death syndrome. *Plant Health Progress*. 2008;10. doi:10.1094/PHP-2008-0222-01-RS.
84. Navi SS, Yang XB. Impact of crop residue and corn-soybean rotation on the survival of *Fusarium virguliforme* a causal agent of sudden death syndrome of soybean. *Journal of Plant Pathology & Microbiology*. 2016 Jan 28;2016. 7:330. doi:10.4172/2157-7471.1000330
85. Navi SS, Yang XB. Seedling inoculation screening technique to identify resistance to soybean sudden death syndrome caused by *Fusarium solani* f. sp. *glycines*. 2005. Pages 252-253. In: Khetarpal RK, Gupta K, Chalam CV, Gour HN, Thakore BBL, Bhargava S, editors. Abstract of Papers Global Conference II on Plant Health-Global Wealth, November 25–29, 2005, Udaipur, India 313 001.
86. Navi SS, Yang XB. Study on penetration by *Fusarium solani* f. sp. *glycines*, causal agent of SDS on soybean. *Phytopathology*. 2004; 94:S74.
87. Navi, SS, Huynh T, Li X, Yang XB. Effects of biocontrol agents in combination with bioAPT microbial carrier as seed treatment on soybean diseases and yields. 2016. Page no 64. In: Khetarpal RK, Mondal KK, Dubey SC, Rao GP, Celia Chalam V, Singh N, et al. editors. IPS 6th International Conference on Plant, Pathogens and People, Feb 23-27, 2016, New Delhi, India. pp 676
88. Nelson KA, Motavalli PP, Stevens WE, Dunn D, Meinhardt CG. Soybean response to preplant and foliar-applied potassium chloride with Strobilurin fungicides. *Agronomy Journal*. 2010;102:1657–63.
89. Njiti VN, Myers O, Schroeder D, Lightfoot DA. Roundup Ready Soybean. *Agronomy Journal*. 2003;95:1140–5.
90. O'Donnell K, Sink S, Scandiani MM, Luque A, Colletto A, Biasoli M, Lenzi L, Salas G, González V, Ploper LD, Formento N. Soybean sudden death syndrome species diversity within North and South America revealed by multilocus genotyping. *Phytopathology*. 2010;100:58–71.

91. Ortiz-Ribbing LM, Eastburn DM. Soybean root systems and sudden death syndrome severity: Taproot and lateral root infection. *Plant Disease*. 2004;88:1011–6.
92. Pennypacker BW. First report of sudden death syndrome caused by *Fusarium solani* f. sp. *glycines* on soybean in Pennsylvania. *Plant Disease*. 1999;83:879.
93. Perez-Brandan C, Arzeno JL, Huidobro J, Conforto C, Grumberg B, Hilton S, et al. The effect of crop sequences on soil microbial, chemical and physical indicators and its relationship with soybean sudden death syndrome (complex of *Fusarium* species). *Spanish Journal of Agricultural Research*. 2013;12:252–64.
94. Ploper D. Síndrome de la muerte súbita: Nueva enfermedad de la soja en el noroeste argentino. *Avance Agroindustrial* Año. 13. 1993;54:5–9.
95. Ploper LD, Yorinori JT, Morel Paiva W. Importancia de las enfermedades de la soja en el Mercosur. In: *Actas Simposio Internacional sobre Soja, XI Congreso de AAPRESID*, Rosario, Argentina, August 2003 Aug 26 (pp. 26-29). [Spanish]
96. Pratt PW, Wrather JA. Soybean disease loss estimates for the southern United States, 1994 to 1996. *Plant Disease*. 1998;82:114–6.
97. Punja ZK, Utkhede RS. Using fungi and yeasts to manage vegetable crop diseases. *TRENDS in Biotechnology*. 2003;21:400–7.
98. Roy KW, Hershman DE, Rupe JC, Abney TS. Sudden death syndrome of soybean. *Plant Disease*. 1997;81:1100–11.
99. Roy KW, Lawrence GW, Hodges HH, McLean KS, Killebrew JF. Sudden death syndrome of soybean: *Fusarium solani* as incitant and relation of *Heterodera glycines* to disease severity. *Phytopathology*. 1989;79:191–7.
100. Roy KW. Sporulation of *Fusarium solani* f. sp. *glycines*, causal agent of sudden death syndrome, on soybeans in the Midwestern and southern United States. *Plant Disease*. 1997;81:566–9.
101. Rupe JC, Gbur EE, Marx DM. Cultivar responses to sudden death syndrome of soybean. *Plant Disease*. 1991;75:47–50.
102. Rupe JC, Hirrel MC, Hershman DE. Sudden death syndrome. In: Sinclair, JB, Backman, PA, editors. *Compendium of Soybean Diseases*. 3rd ed. APS Press, St. Paul, MN, USA; 1989. pp. 84–5.

103. Rupe JC, Robbins RT, Becton CM, Sabbe WA, Gbur EE. Vertical and temporal distribution of *Fusarium solani* and *Heterodera glycines* in fields with sudden death syndrome of soybean. *Soil Biology and Biochemistry*. 1999;31:245–51.
104. Rupe JC, Robbins RT, Gbur EE. Effect of crop rotation on soil population densities of *Fusarium solani* and *Heterodera glycines* and on the development of sudden death syndrome of soybean. *Crop Protection*. 1997;16:575–80.
105. Rupe JC, Sabbe WE, Robbins RT, Gbur EE. Soil and plant factors associated with sudden death syndrome of soybean. *Journal of Production Agriculture*. 1993;6:218–21.
106. Samuels GJ, Hebbar PK. Developing Trichoderma-based products for application in agriculture. 2015. Pages 7-34, In: *Trichoderma identification and agricultural applications*. The American Phytopathology, 3340 Pilot knob road, St. Paul, Minnesota, U.S.A. Edited by Gary J. Samuels and Prakash K. Hebbar
107. Sanogo S, Yang XB, Lundeen P. Field response of glyphosate-tolerant soybean to herbicides and sudden death syndrome. *Plant Disease*. 2001;85:773–9.
108. Sanogo S, Yang XB, Scherm H. Effects of herbicides on *Fusarium solani* f. sp. *glycines* and development of sudden death syndrome in glyphosate-tolerant soybean. *Phytopathology*. 2000;90:57–66.
109. Sanogo S, Yang XB. Relation of sand content, pH, and potassium and phosphorus nutrition to the development of sudden death syndrome in soybean. *Canadian Journal of Plant Pathology*. 2001;23:174–80.
110. Scandiani M, Ruberti D, Pioli R, Luque A, Giorda L. First report of Koch's postulates completion of sudden death syndrome of soybean in Argentina. *Plant Disease*. 2003;87:447.
111. Scandiani MM, Ruberti DS, Giorda LM, Pioli RN, Luque AG, Bottai H, et al. Comparison of inoculation methods for characterizing relative aggressiveness of two soybean sudden-death syndrome pathogens, *Fusarium virguliforme* and *F. tucumaniae*. *Tropical Plant Pathology*. 2011;36:133–40.
112. Scherm H, Yang XB, Lundeen P. Soil variables associated with sudden death syndrome in soybean fields in Iowa. *Plant Disease*. 1998;82:1152–7.
113. Scherm H, Yang XB. Development of sudden death syndrome of soybean in relation to soil temperature and soil water matric potential. *Phytopathology*. 1996;86:642–9.

114. Scherm H, Yang XB. Risk assessment for sudden death syndrome of soybean in the north-central United States. *Agricultural Systems*. 1999;59:301–10.
115. Schirmböck M, Lorito M, Wang YL, Hayes CK, Arisan-Atac I, Scala F, et al. Parallel formation and synergism of hydrolytic enzymes and peptaibol antibiotics, molecular mechanisms involved in the antagonistic action of *Trichoderma harzianum* against phytopathogenic fungi. *Applied and Environmental Microbiology*. 1994;60:4364–70.
116. Sciumbato GL. Soybean disease loss estimates for the southern United States during 1988-1991. *Plant Disease*. 1993;77:954–6.
117. Seyb A, Xing L, Vyn TJ, Seo J, Abney S, Westphal A. Tillage system effects on sudden death syndrome, *Heterodera glycines*, and soybean yield in a Mollisol. In *Phytopathology* 2007 Jul 1 (Vol. 97, No. 7, pp. S106-S107). 3340 Pilot Knob Road, St Paul, MN 55121 USA: American Phytopathological Society.
118. Sikora EJ, Delaney D. Sudden death syndrome of soybeans In: *Plant Pathology Series, timely information on agriculture and natural resources*, Alabama Cooperative Extension System, 2004; page 572. <https://sites.aces.edu/group/timelyinfo/Documents/pp572.pdf>
119. Singh R, Price T, Padgett B, Burks T. First report of sudden death syndrome of soybean caused by *Fusarium virguliforme* in Louisiana. *Plant Health Progress*. 2015. *Plant Health Progress* doi:10.1094/PHP-BR-15-0031
120. Singh SD, De Milliano WA. Production of normal panicles by sorghum plants systemically infected by downy mildew in Zimbabwe. *Plant Disease*. 1989;73:1020.
121. Singh SD, King SB. Recovery resistance to downy mildew in pearl millet. *Plant Disease*. 1988;72:425–8.
122. Sivan A, Chet I. Biological control of *Fusarium* spp. in cotton, wheat and muskmelon by *Trichoderma harzianum*. *Journal of Phytopathology*. 1986;116:39–47.
123. Swoboda CM. Effect of agronomic practices on sudden death syndrome of soybean in Iowa. 2010. Graduate Theses and Dissertations, Paper 11559.
124. Tande C, Hadi B, Chowdhury R, Subramanian S, Byamukama E. First report of sudden death syndrome of soybean caused by *Fusarium virguliforme* in South Dakota. *Plant Disease*. 2014;98:1012.

125. Tewoldemedhin YT, Lamprecht SC, Geldenhuys JJ, Kloppers FJ. First report of soybean sudden death syndrome caused by *Fusarium virguliforme* in South Africa. *Plant Disease*. 2014;98:569.
126. Verma M, Brar SK, Tyagi RD, Surampalli RY, Valero JR. Antagonistic fungi, *Trichoderma* spp.: panoply of biological control. *Biochemical Engineering Journal*. 2007;37:1–20.
127. Vick CM, Bond JP, Chong SK, Russin JS. Response of soybean sudden death syndrome to tillage and cultivar. *Canadian Journal of Plant Pathology*. 2006;28:77–83.
128. Vick CM, Chong SK, Bond JP, Russin JS. Response of soybean sudden death syndrome to subsoil tillage. *Plant Disease*. 2003;87:629–32.
129. Warner A, Arnao E, Fakhoury A. The use of fungal biological control agents to manage SDS, charcoal rot, and damping off in soybean. 2016. APS annual meeting, July 30 - Aug 3, Tampa, Florida, U.S.A.
<http://www.apsnet.org/meetings/annual/abstracts/pages/abstractdetail.aspx?MID=408>
130. Weems JD, Haudenshield JS, Bond JP, Hartman GL, Ames KA, Bradley CA. Effect of fungicide seed treatments on *Fusarium virguliforme* infection of soybean and development of sudden death syndrome. *Canadian Journal of Plant Pathology*. 2015;37:435–447.
131. Weems JD, Zhang GR, Ames KA, Haudenshield JS, Hartman GL, Bond JP, et al. Effect of fungicide seed treatments on *Fusarium virguliforme* and sudden death syndrome of soybean. In American Pathological Society North Central Division Annual Meeting 2010 Jun 8.
132. Wrather JA, Anderson TR, Arsyad DM, Gai J, Ploper LD, Porta-Puglia A, Ram HH, Yorinori JT. Soybean disease loss estimates for the top 10 soybean producing countries in 1994. *Plant Disease*. 1997;81:107–10.
133. Wrather JA, Anderson TR, Arsyad DM, Tan Y, Ploper LD, Porta-Puglia A, et al. Soybean disease loss estimates for the top ten soybean-producing countries in 1998. *Canadian Journal of Plant Pathology*. 2001;23:115–21.
134. Wrather JA, Chambers AY, Fox JA, Moore WF, Sciumbato GL. Soybean disease loss estimates for the southern United States, 1974 to 1994. *Plant Disease*. 1995;79:1076–99.
135. Wrather JA, Kendig SR, Anand SC, Niblack TL, Smith GS. Effects of tillage, cultivar, and planting date on percentage of soybean leaves with symptoms of sudden death syndrome. *Plant Disease*. 1995a;79:560–2.

136. Wrather JA, Koenning SR, Anderson TR. Effect of diseases on soybean yields in the United States and Ontario (1999–2002). *Plant Health Progress*. 2003;1. doi:10.1094/PHP-2003-0325-01-RV.
137. Wrather JA, Koenning SR. Effects of diseases on soybean yields in the United States 1996 to 2007. *Plant Health Progress*. 2009;10. doi:10.1094/PHP-2009-0401-01-RS.
138. Wrather JA, Koenning SR. Estimates of disease effects on soybean yields in the United States 2003 to 2005. *Journal of Nematology*. 2006;38:173.
139. Wrather JA, Koenning SR. Soybean disease loss estimates for the United States, 1996–2010. Delta Research Center: agriculture experiment station. University of Missouri, College of agriculture, food and natural resources. Food and Natural Resources. 2011.
140. Wrather JA, Sciumbato GL. Soybean disease loss estimates for the southern United States during 1992 and 1993. *Plant Disease (USA)*. 1995.
141. Wrather JA, Stienstra WC, Koenning SR. Soybean disease loss estimates for the United States from 1996 to 1998. *Canadian Journal of Plant Pathology*. 2001a;23:122–31.
142. Xing L, Westphal A. Effects of crop rotation of soybean with corn on severity of sudden death syndrome and population densities of *Heterodera glycines* in naturally infested soil. *Field Crops Research*. 2009;112:107–17.
143. Xing L, Westphal A. Interaction of *Fusarium solani* f. sp. *glycines* and *Heterodera glycines* in sudden death syndrome of soybean. *Phytopathology*. 2006;96:763–70.
144. Xing L, Westphal A. Synergism in the interaction of *Fusarium virguliforme* with *Heterodera glycines* in sudden death syndrome of soybean. *Journal of Plant Diseases and Protection*. 2013;120:209–17.
145. Yamanaka N, Fuentes FH, Gilli JR, Watanabe S, Harada K, Ban T, et al. Identification of quantitative trait loci for resistance against soybean sudden death syndrome caused by *Fusarium tucumaniae*. *Pesquisa Agropecuária Brasileira*. 2006;41:1385–91.
146. Yang S, Li X, Chen C, Kyveryga P, Yang XB. Assessing field specific risk of soybean sudden death syndrome using satellite imagery in Iowa. *Phytopathology*. 2016;106:842–53.
147. Yang X. Occurrence of soybean sudden death syndrome after adoption of Roundup Ready technology in North America. *Phytopathology*. 2012;102:140.

148. Yang XB, Lundeen P, Uphoff M. An efficient method for field inoculation of soybean sudden death syndrome caused by *Fusarium solani* f. sp. *glycines*. *Phytopathology*. 2001;91:S97.
149. Yang XB, Navi SS. Strategies for management of soybean sudden death syndrome and white mold. Pages 105–112. in B. Pringnitz (Ed.), *Proceedings of the 18th Annual Integrated Crop Management Conference*. 2006. Ames, IA: Iowa State University.
150. Yang XB, Rizvi, SSA. First report of sudden death syndrome of soybean in Iowa. *Plant Disease*. 1994;78:830.
151. Yang, XB. Navi, SS. Good harvest in corn should help manage soybean SDS. Iowa State University, *Integrated Crop Management News*. 2010; September 10.
152. Yorinori JT. Situação atual das doenças potenciais no Cone Sul. In Congresso Brasileiro de Soja 2002 (Vol. 2, pp. 171–186).
153. Yorinori JT. Situação atual das enfermidades de soja na Bolívia, Brasil e Paraguai. *Proceedings, Mercosoja*. 1999;99:21–5.
154. Zaccaron ML, Yang XB, Navi SS. Foliar symptoms expression and early infection of soybean sudden death syndrome. *Phytopathology*. 2010;100:S144.
155. Zaworski ER. Effects of IleVO® on soybean sudden death syndrome and soybean cyst nematode. Graduate Theses and Dissertations. 2014. Paper 14261. Iowa State University, Ames, IA.
156. Zhang J, Singh A, Mueller DS, Singh AK. Genome-wide association and epistasis studies unravel the genetic architecture of sudden death syndrome resistance in soybean. *The Plant Journal*. 2015;84:1124–36.
157. Ziems AD, Giesler LJ, Yuen GY. First report of sudden death syndrome of soybean caused by *Fusarium solani* f. sp. *glycines* in Nebraska. *Plant Disease*. 2006;90:101.




Figure 2: Yield and economic losses due to soybean sudden death syndrome (SDS) for the U.S.A. soybean production from 1988 to 2010. [Data source for yield loss: 21, 35, 53, 96, 116, 132–134, 136–141 and for soybean prices: USDA Economic Research Service www.ers.usda.gov/]




Figure 3: Field symptoms of soybean sudden death syndrome caused by *Fusarium virguliforme* in seedling stage; (A) chlorotic spots, (B) chlorosis, (C) stunting, (D) necrosis and puckering and (E) discolored basal stem, tap roots and root hairs of infected plants (left) and uninfected plant with healthy roots and leaves (right) and in reproductive growth stages; (F) chlorotic spots and chlorosis, (G) necrosis, (H) severe necrosis and puckering, (I) loss of leaf lamina (J) defoliation with leaf petioles intact on stems and (K) *F. virguliforme* growth on taproot.




Figure 4: Effective and ineffective colonization zones of *Fusarium virguliforme* in the section of a soybean taproot [83].




Figure 5: Disease cycle of sudden death syndrome of soybean caused by *Fusarium virguliforme*.




Figure 6: Effects of seed treatments with either HeadsUp alone or HeadsUp plus commercial seed treatment products on SDS reduction as indicated by yield increase. Trials were conducted in fields infested with SDS fungus in the U.S. Midwestern regions.



1
2 Figure 7: Greenhouse screening technique for: 1. efficacy tests of biocontrol agents or fungicides seed treatment and 2. identification of sources of
3 resistance to soybean sudden death syndrome; (A) spread inoculum at 1% of the potting mixture in plastic tub, (B) Mix the inoculum with potting
4 mixture, (C) Fill-up the cups, (D) Mark planting spots with hand held wooden planter (WP), (E) Place 5-7 seeds in marked spots and press the seed
5 to uniform depth of 2-cm with WP, (F) Cover the seed with potting mixture and (G) Compact it with hand held wooden compactor (WC). Water twice
6 daily to maintain enough moisture in cups, and F. Record stand counts and SDS counts 15, 30 and 45 days after planting.